

Topic:- DU_J19_MA_MATHS

1) The order of Sylow subgroups of a finite group G of order 56 are [Question ID = 24519]

1. 2 and 28 [Option ID = 38076]
2. 7 and 8 [Option ID = 38074]
3. 8 and 14 [Option ID = 38077]
4. 4 and 14 [Option ID = 38075]

Correct Answer :-

- 7 and 8 [Option ID = 38074]

2) The remainder when 5^{2019} is divided by 11 is [Question ID = 24520]

1. 6 [Option ID = 38080]
2. 9 [Option ID = 38081]
3. 1 [Option ID = 38078]
4. 4 [Option ID = 38079]

Correct Answer :-

- 9 [Option ID = 38081]

3) The smallest positive integer n , which leaves remainders 2,3 and 4 when divided by 5,7 and 11 respectively, is [Question ID = 24521]

1. 751 [Option ID = 38083]
2. 1136 [Option ID = 38085]
3. 176 [Option ID = 38082]
4. 367 [Option ID = 38084]

Correct Answer :-

- 367 [Option ID = 38084]

4) Suppose that the equation $x^2 \cdot a \cdot x = a^{-1}$ is solvable for a in a group G . Then, there exists b in G such that [Question ID = 24515]

1. $a = b^3$ [Option ID = 38059]
2. $a = b^5$ [Option ID = 38061]
3. $a = b^4$ [Option ID = 38060]
4. $a = b^2$ [Option ID = 38058]

Correct Answer :-

- $a = b^3$ [Option ID = 38059]

5) Consider the following statements:

- (i) Every metric space is totally bounded.
- (ii) A totally bounded metric space is bounded.

Then

[Question ID = 24536]

1. neither (i) nor (ii) is true [Option ID = 38145]
2. only (ii) is true [Option ID = 38143]
3. only (i) is true [Option ID = 38142]
4. both (i) and (ii) are true [Option ID = 38144]

Correct Answer :-

- only (ii) is true [Option ID = 38143]

6)

Consider the following statements:

- (i) Every minimal generating set of a vector space is a basis.
- (ii) Every maximal linearly independent subset of a vector space is a basis.
- (iii) Every vector space admits a basis.

Then

[Question ID = 24510]

- 1. all of (i), (ii) and (iii) are true [Option ID = 38041]
- 2. only (i) and (ii) are true [Option ID = 38038]
- 3. only (i) and (iii) are true [Option ID = 38040]
- 4. only (ii) and (iii) are true [Option ID = 38039]

Correct Answer :-

- all of (i), (ii) and (iii) are true [Option ID = 38041]

- 7) The differential equation of a family of parabolas with foci at origin and axis along x -axis is

[Question ID = 24506]

- 1. $y\left(\frac{dy}{dx}\right)^2 + 2x^2\frac{dy}{dx} + y = 0$ [Option ID = 38023]
- 2. $y\left(\frac{dy}{dx}\right)^2 + 2x\frac{dy}{dx} - y = 0$ [Option ID = 38024]
- 3. $y\left(\frac{dy}{dx}\right)^2 + 2x\frac{dy}{dx} + y = 0$ [Option ID = 38025]
- 4. $y^2\left(\frac{dy}{dx}\right)^2 + 2x\frac{dy}{dx} - y^2 = 0$ [Option ID = 38022]

Correct Answer :-

- $y\left(\frac{dy}{dx}\right)^2 + 2x\frac{dy}{dx} - y = 0$ [Option ID = 38024]

- 8) Number of iterations required to solve $x^3 + 4x^2 - 10 = 0$ using bisection method with accuracy 10^{-3} (with initial bracket $[1, 2]$) are

[Question ID = 24495]

- 1. 7 [Option ID = 37978]
- 2. 12 [Option ID = 37981]
- 3. 10 [Option ID = 37980]
- 4. 8 [Option ID = 37979]

Correct Answer :-

- 10 [Option ID = 37980]

- 9) Let $P_2(t)$ denote the set of all polynomials over \mathbb{R} of degree at most 2. With respect to the inner product

$$\langle p, q \rangle = \int_{-1}^1 p(t)q(t)dt,$$

the set of vectors $\{1, t, t^2 - \frac{1}{3}\}$ is

[Question ID = 24513]

- 1. not a linearly independent set [Option ID = 38053]
- 2. orthogonal basis of $P_2(t)$ [Option ID = 38050]
- 3. basis of $P_2(t)$ but not orthogonal [Option ID = 38052]

4. orthogonal but not a basis of $P_2(t)$ [Option ID = 38051]

Correct Answer :-

- orthogonal basis of $P_2(t)$ [Option ID = 38050]

10) A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be periodic if there exists $p > 0$ such that $f(x + p) = f(x)$, for all $x \in \mathbb{R}$. If f is a continuous periodic function on \mathbb{R} , then

[Question ID = 24543]

1. f^2 is unbounded [Option ID = 38173]
2. $|f|$ is unbounded [Option ID = 38170]
3. $|f|$ is not uniformly continuous [Option ID = 38172]
4. f^2 is uniformly continuous and bounded on \mathbb{R} [Option ID = 38171]

Correct Answer :-

- f^2 is uniformly continuous and bounded on \mathbb{R} [Option ID = 38171]

11) Consider the following statements:

- (i) Every separable metric space is compact.
- (ii) Every compact metric space is separable.

Then

[Question ID = 24534]

1. only (i) is true [Option ID = 38134]
2. only (ii) is true [Option ID = 38135]
3. both (i) and (ii) are true [Option ID = 38136]
4. neither (i) nor (ii) is true [Option ID = 38137]

Correct Answer :-

- only (ii) is true [Option ID = 38135]

12) The partial differential equation $x^3 u_{xx} - (y^2 - 1)u_{yy} = u_x$ is

[Question ID = 24502]

1. parabolic in $\{(x, y) \mid y < 0\}$ [Option ID = 38006]
2. elliptic in \mathbb{R}^2 [Option ID = 38008]
3. hyperbolic in $\{(x, y) \mid x > 0\}$ [Option ID = 38007]
4. parabolic in $\{(x, y) \mid y > 0\}$ [Option ID = 38009]

Correct Answer :-

13) Consider the following statements

- (i) $\mathbb{Z}[x]$ is a principal ideal domain.
- (ii) If R is a principal ideal domain, then every subring of R containing 1 is also a principal ideal domain.

Then

[Question ID = 24522]

1. only (i) is true [Option ID = 38086]

2. both (i) and (ii) are true [Option ID = 38088]
3. only (ii) is true [Option ID = 38087]
4. neither (i) nor (ii) is true [Option ID = 38089]

Correct Answer :-

- neither (i) nor (ii) is true [Option ID = 38089]

14) Let $N \neq \{e\}$ be a normal subgroup of a non-abelian group G such that $N \cap G' = \{e\}$, where G' is the commutator subgroup of G . Then

[Question ID = 24517]

1. None of these [Option ID = 38069]
2. N is not abelian [Option ID = 38067]
3. $N \subseteq Z(G)$, the centre of G [Option ID = 38068]
4. G/N is abelian [Option ID = 38066]

Correct Answer :-

- $N \subseteq Z(G)$, the centre of G [Option ID = 38068]

15) Let $f(t) = t^2 e^t \log t$; $1 \leq t \leq 3$. Then there exists some $c \in (1, 3)$ such that $\int_1^3 f(t) dt$ is equal to

[Question ID = 24525]

1. $\frac{1}{3} e^c \log c^{26}$ [Option ID = 38098]
2. $c^2 e^c \log 3$ [Option ID = 38101]
3. $2^2 c^2 \log c$ [Option ID = 38099]
4. $26 e^c \log c$ [Option ID = 38100]

Correct Answer :-

- $\frac{1}{3} e^c \log c^{26}$ [Option ID = 38098]

16) For two ideals I and J of a commutative ring R define $(I : J) = \{r \in R \mid rI \subseteq J\}$. Then for the ring \mathbb{Z} of integers what is $(8\mathbb{Z} : 12\mathbb{Z})$

[Question ID = 24523]

1. $4\mathbb{Z}$ [Option ID = 38093]
2. \mathbb{Z} [Option ID = 38090]
3. $2\mathbb{Z}$ [Option ID = 38091]
4. $3\mathbb{Z}$ [Option ID = 38092]

Correct Answer :-

- $3\mathbb{Z}$ [Option ID = 38092]

17) Consider the set \mathbb{R}^2 with metric defined by

$$d(x, y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}; \quad x = (x_1, x_2), \quad y = (y_1, y_2).$$

Then which of the following set is not connected

[Question ID = 24535]

1. $\{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$ [Option ID = 38138]
2. $\{(x, y) \in \mathbb{R}^2 \mid x^2 - y^2 = 1\}$ [Option ID = 38141]
3. $\{(x, y) \in \mathbb{R}^2 \mid \frac{x^2}{4} + \frac{y^2}{9} = 1\}$ [Option ID = 38140]
4. $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ [Option ID = 38139]

Correct Answer :-

- $\{(x, y) \in \mathbb{R}^2 \mid x^2 - y^2 = 1\}$ [Option ID = 38141]

18) Let $f(x) = \lim_{n \rightarrow \infty} \frac{n^x - n^{-x}}{n^x + n^{-x}}, x \in \mathbb{R}$. Then

[Question ID = 24542]

1. f is continuous at $(1, \infty)$ [Option ID = 38169]
2. f is not differentiable at $x = 1$ [Option ID = 38168]
3. f is not continuous at $x = -1$ [Option ID = 38167]
4. f is continuous at $x = 0$ [Option ID = 38166]

Correct Answer :-

- f is continuous at $(1, \infty)$ [Option ID = 38169]

19) For $x \in [-1, 1]$, let

$$f(x) = \begin{cases} x \operatorname{sgn}(\sin \frac{1}{x}), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0, \end{cases}$$

where sgn denotes the signum function. Then

[Question ID = 24526]

1. f is continuous on $[-1, 1]$ [Option ID = 38104]
2. f is not differentiable at any point of $[-1, 1]$ [Option ID = 38103]
3. f is Riemann integrable on $[-1, 1]$ [Option ID = 38102]
4. the set of points of discontinuity of f in $[-1, 1]$ is finite [Option ID = 38105]

Correct Answer :-

- f is Riemann integrable on $[-1, 1]$ [Option ID = 38102]

20) The integral surface of the partial differential equation $p^2 + q^2 = 2$ which passess through $x = 0, z = y$ is

[Question ID = 24503]

1. $x^2 + y^2 + z^2 = 1$ [Option ID = 38013]
2. $z = y \pm x$ [Option ID = 38010]
3. $z^2 = x \pm y^2$ [Option ID = 38011]
4. $z^3 = x \pm y$ [Option ID = 38012]

Correct Answer :-

- $z = y \pm x$ [Option ID = 38010]

21) Does the sequence $a_n = n^2 \cos\left(\frac{2}{n^2} + \frac{\pi}{2}\right)$ has a limit?

[Question ID = 24529]

1. No, it oscillates [Option ID = 38115]
2. No, it diverges [Option ID = 38114]
3. Yes, -2 is the limit [Option ID = 38117]
4. Yes, -1 is the limit [Option ID = 38116]

Correct Answer :-

- Yes, -2 is the limit [Option ID = 38117]

22)

The orthogonal trajectory of the family of curves $ay^2 = x^3$, where a is an arbitrary constant, is

[Question ID = 26021]

1. $3y^2 + 2x^2 = \text{constant}$ [Option ID = 44082]
2. $2y^2 - 3x^2 = \text{constant}$ [Option ID = 44080]
3. $3y^2 - 2x^2 = \text{constant}$ [Option ID = 44079]
4. $2y^2 + 3x^2 = \text{constant}$ [Option ID = 44081]

Correct Answer :-

- $3y^2 + 2x^2 = \text{constant}$ [Option ID = 44082]

23) The integral surface of the linear partial differential equation

$$xp + yq = z$$

which contains the circle defined by $x^2 + y^2 + z^2 = 4$, $x + y + z = 2$, is

[Question ID = 24504]

1. $\frac{x}{y} + \frac{z}{x} + \frac{y}{z} + 1 = 0$ [Option ID = 38015]
2. $xy + xz + yz = 0$ [Option ID = 38016]
3. $xy^2 + xz^2 = 0$ [Option ID = 38014]
4. $xyz = 1$ [Option ID = 38017]

Correct Answer :-

- $xy + xz + yz = 0$ [Option ID = 38016]

24) Initial estimate for the root of the equation $f(x) = 0$ is $x_0 = 2$ and $f(2) = 4$. The tangent line to $f(x)$ at $x_0 = 2$ makes an angle of 42° with the x axis. The next estimate of the root by Newton-Raphson method is approximately

[Question ID = 24499]

1. 2.0102 [Option ID = 37995]
2. 4.4424 [Option ID = 37997]
3. 0.2412 [Option ID = 37994]
4. -2.4424 [Option ID = 37996]

Correct Answer :-

- -2.4424 [Option ID = 37996]

25) The numerical scheme using the first three terms of the Taylor series for solving the differential equation

$$\frac{dy}{dx} + y = e^{-3x}, \quad y(0) = 5,$$

with $h = x_{i+1} - x_i$, is given by

[Question ID = 24497]

1. $y_{i+1} = y_i + h(e^{-3x_i} - y_i) + \frac{h^2}{2}(-3e^{-3x_i} - y_i)$ [Option ID = 37988]
2. $y_{i+1} = y_i + h(e^{-3x_i} - y_i) + \frac{h^2}{2}(-4e^{-3x_i} + y_i)$ [Option ID = 37987]
3. $y_{i+1} = y_i - h(e^{-3x_i} - y_i) + \frac{h^2}{2}(y_i - e^{-3x_i})$ [Option ID = 37989]
4. $y_{i+1} = y_i + h(e^{-3x_i} - y_i) + \frac{h^2}{2}y_i$ [Option ID = 37986]

Correct Answer :-

- $y_{i+1} = y_i + h(e^{-3x_i} - y_i) + \frac{h^2}{2}(-4e^{-3x_i} + y_i)$ [Option ID = 37987]

26) Let $X = \mathbb{C}^n$, $0 < p < 1$ and $q = 1/p$. For $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ in X define

$$d_p(x, y) = \left(\sum_{i=1}^n |x_i - y_i|^p \right)^{1/p}$$

and

$$d_q(x, y) = \left(\sum_{i=1}^n |x_i - y_i|^q \right)^{1/q}.$$

Then

[Question ID = 24533]

1. neither $d_p(x, y)$ nor $d_q(x, y)$ is a metric on X [Option ID = 38133]
2. both $d_p(x, y)$ and $d_q(x, y)$ are metrics on X [Option ID = 38130]
3. only $d_q(x, y)$ is a metric on X [Option ID = 38132]
4. only $d_p(x, y)$ is a metric on X [Option ID = 38131]

Correct Answer :-

- only $d_q(x, y)$ is a metric on X [Option ID = 38132]

27) Let $f(x) = x \sin x$, $x \in \mathbb{R}$. Then $|f|$ is

[Question ID = 26030]

1. differentiable at $x = \pi$ [Option ID = 44117]
2. differentiable at $x = 0$ [Option ID = 44115]
3. uniformly continuous on \mathbb{R} [Option ID = 44118]
4. differentiable at $x = -\pi$ [Option ID = 44116]

Correct Answer :-

- differentiable at $x = 0$ [Option ID = 44115]

28) Which of the following function f is not uniformly continuous on \mathbb{R}

[Question ID = 24541]

1. $f(x) = x + \sin x$ [Option ID = 38163]
2. $f(x) = x + \sin^3 x$ [Option ID = 38165]
3. [Option ID = 38164]
4. [Option ID = 38162]

Correct Answer :-

- [Option ID = 38164]

29)
[Question ID = 24512]

1. X and Z [Option ID = 38047]
2. Y and Z [Option ID = 38049]
3. W and Y [Option ID = 38046]
4. W and X [Option ID = 38048]

Correct Answer :-

- X and Z [Option ID = 38047]

30)
[Question ID = 24509]

1. [Option ID = 38034]
2. [Option ID = 38037]
3. [Option ID = 38036]
4. [Option ID = 38035]

Correct Answer :-

- [Option ID = 38035]

31)
[Question ID = 24539]

1. [Option ID = 38156]
2. [Option ID = 38155]
3. [Option ID = 38157]
4. [Option ID = 38154]

Correct Answer :-

- [Option ID = 38155]

32)
[Question ID = 24507]

1. [Option ID = 38027]
2. [Option ID = 38026]
3. [Option ID = 38028]
4. [Option ID = 38029]

Correct Answer :-

- [Option ID = 38029]

33)
[Question ID = 24511]

1. only (ii) and (iii) are true [Option ID = 38044]
2. only (ii), (iii) and (iv) are true [Option ID = 38043]
3. only (i) and (ii) are true [Option ID = 38045]
4. only (i), (ii) and (iii) are true [Option ID = 38042]

Correct Answer :-

- only (i), (ii) and (iii) are true [Option ID = 38042]

34)

[Question ID = 24500]

1. [Option ID = 38000]
2. [Option ID = 37998]
3. [Option ID = 38001]
4. [Option ID = 37999]

Correct Answer :-

- [Option ID = 37998]

35)

[Question ID = 24496]

1. [Option ID = 37983]
2. [Option ID = 37982]
3. [Option ID = 37985]
4. [Option ID = 37984]

Correct Answer :-

- [Option ID = 37982]

36)

[Question ID = 24514]

1. S is one-one but T is not [Option ID = 38055]
2. T is one-one but S is not [Option ID = 38054]
3. Both S and T are one-one [Option ID = 38056]
4. Neither S nor T is one-one [Option ID = 38057]

Correct Answer :-

- Both S and T are one-one [Option ID = 38056]

37)

[Question ID = 24501]

1. [Option ID = 38002]
2. [Option ID = 38004]
3. [Option ID = 38005]
4. [Option ID = 38003]

Correct Answer :-

- [Option ID = 38005]

38)

[Question ID = 24518]

1. both c and d are even [Option ID = 38072]
2. both c and d are odd [Option ID = 38073]
3. c is even and d is odd [Option ID = 38071]
4. c is odd and d is even [Option ID = 38070]

Correct Answer :-

- c is odd and d is even [Option ID = 38070]

39)

[Question ID = 24524]

1. only (ii) and (iii) are true [Option ID = 38095]
2. only (i) and (iii) are true [Option ID = 38096]
3. only (i) and (ii) are true [Option ID = 38094]
4. all of (i), (ii) and (iii) are true [Option ID = 38097]

Correct Answer :-

- only (i) and (ii) are true [Option ID = 38094]

40)
[Question ID = 24537]

1. [Option ID = 38146]
2. [Option ID = 38147]
3. [Option ID = 38148]
4. [Option ID = 38149]

Correct Answer :-

- [Option ID = 38147]

41)
[Question ID = 24527]

1. [Option ID = 38109]
2. [Option ID = 38108]
3. [Option ID = 38106]
4. [Option ID = 38107]

Correct Answer :-

- [Option ID = 38108]

42)
[Question ID = 24528]

1. [Option ID = 38111]
2. [Option ID = 38113]
3. [Option ID = 38112]
4. [Option ID = 38110]

Correct Answer :-

- [Option ID = 38112]

43)
[Question ID = 24530]

1. all of (i), (ii) and (iii) are true [Option ID = 38121]
2. only (ii) is true [Option ID = 38119]
3. only (i) and (ii) are true [Option ID = 38118]
4. only (ii) and (iii) are true [Option ID = 38120]

Correct Answer :-

- all of (i), (ii) and (iii) are true [Option ID = 38121]

44)
[Question ID = 24498]

1. 0.0996 [Option ID = 37991]
2. 0.0876 [Option ID = 37990]
3. 0.0745 [Option ID = 37992]
4. 0.0912 [Option ID = 37993]

Correct Answer :-

- 0.0996 [Option ID = 37991]

45)
[Question ID = 24531]

1. converges for all values of p [Option ID = 38124]
2. converges for $p > 0$, diverges for $p \leq 0$ [Option ID = 38122]
3. does not converges for any value of p [Option ID = 38125]
4. converges for $p > 1$, diverges for $p \leq 1$ [Option ID = 38123]

Correct Answer :-

- converges for $p > 0$, diverges for $p \leq 0$ [Option ID = 38122]

46)
[Question ID = 26022]

1. [Option ID = 44086]
2. [Option ID = 44083]
3. [Option ID = 44084]
4. [Option ID = 44085]

Correct Answer :-

- [Option ID = 44083]

47)

[Question ID = 24538]

1. None of these [Option ID = 38153]
2. [Option ID = 38152]
3. [Option ID = 38151]
4. [Option ID = 38150]

Correct Answer :-

- [Option ID = 38150]

48)

[Question ID = 24532]

1. All of X, Y and Z [Option ID = 38126]
2. Only Y and Z [Option ID = 38127]
3. Only X and Z [Option ID = 38128]
4. Only Z [Option ID = 38129]

Correct Answer :-

- Only Z [Option ID = 38129]

49)

[Question ID = 24540]

1. [Option ID = 38161]
2. [Option ID = 38159]
3. [Option ID = 38160]
4. [Option ID = 38158]

Correct Answer :-

50) Let K be any subgroup of a group G and H be the only subgroup of order m in G . Which of the following is not true?

[Question ID = 24516]

1. H is a normal subgroup of G [Option ID = 38062]
2. $G = N(H)$, where $N(H)$ is the normalizer of H in G . [Option ID = 38065]
3. $ab \in H$ implies that $ba \in H$ [Option ID = 38064]
4. HK is not a subgroup of G [Option ID = 38063]

Correct Answer :-

- HK is not a subgroup of G [Option ID = 38063]